

## Osteogenesis Imperfecta (OI) and Dentistry

Last updated: 08 November 2024

**Osteogenesis Imperfecta (OI)**, also known as "brittle bone disease," is a genetic disorder characterised by fragile bones that break easily. It results from mutations in the genes responsible for producing collagen, a protein that provides strength and structure to bones and other connective tissues. The severity of OI can vary, with some individuals experiencing frequent fractures, while others may have milder symptoms.

When it comes to **dentistry** and OI, several challenges arise due to the systemic nature of the disease, particularly affecting the bones and connective tissues, which also include the teeth. Here are key dental considerations and implications for patients with OI:

### 1. Dentinogenesis Imperfecta (DI):

One of the most notable dental concerns in OI is **dentinogenesis imperfecta**, a disorder of tooth development that often accompanies OI. It affects the dentin (the layer beneath the enamel) and leads to:

- **Discoloured teeth:** Teeth may appear blue, grey, or yellowish due to the translucency of the enamel over discoloured dentin.
- **Weakened tooth structure:** The dentin is less mineralised, which makes the teeth more prone to wear, breakage, and cavities.
- **Chipped or cracked teeth:** Teeth can be brittle and prone to fractures, even with minor trauma or normal chewing forces.

### 2. Increased Risk of Dental Problems:

Due to the fragility of both the teeth and bones in individuals with OI, there are several dental issues to consider:

- **Tooth loss:** Because of the brittle nature of teeth and dentinogenesis imperfecta, tooth loss can occur more easily.
- **Difficulty with oral hygiene:** Due to the potential for pain and fractures, patients with OI may find it challenging to maintain regular brushing and flossing routines.
- **Increased susceptibility to dental caries (cavities):** The weakened enamel and dentin may be more susceptible to decay.

### 3. Sensitivity to Trauma:

Individuals with OI are more susceptible to fractures and injury from trauma. This includes:

- **Avoidance of invasive dental procedures:** Invasive treatments, such as extractions or root canals, may require extra care due to the risk of fractures in both the teeth and surrounding bone.
- **Non-invasive techniques:** Dentists may opt for conservative treatments that minimise the risk of further trauma. For example, crowns and veneers can be used to protect weakened teeth, and dental sealants can help prevent cavities.

#### 4. Orthodontic Considerations:

Orthodontic treatment in patients with OI is complex due to several factors:

- **Bone fragility:** The bone may be too fragile to support traditional orthodontic appliances, such as braces.
- **Delayed tooth eruption:** The development of teeth may be delayed or irregular, which can complicate orthodontic planning.
- **Conservative management:** When orthodontic treatment is necessary, it may be adjusted to avoid excessive force on the teeth and bone.

#### 5. Anaesthesia and Pain Management:

- **Anaesthesia:** Patients with OI may have different responses to anaesthesia due to potential issues with connective tissue and bone density. Local anaesthesia is generally safe, but special care must be taken when performing dental procedures.
- **Pain management:** Individuals with OI may experience more intense pain in the teeth and gums due to the fragile nature of their bones and dentin, so effective pain control is essential.

#### 6. Preventive Care:

Due to the increased risk of dental problems, preventive care is critical for individuals with OI:

- **Frequent dental checkups:** Regular dental visits are essential to monitor the health of the teeth and gums and catch any issues early.
- **Fluoride treatments:** Fluoride can help strengthen enamel and reduce the risk of cavities.
- **Customised oral hygiene products:** soft toothbrushes, non-abrasive toothpaste, and gentle flossing techniques can help patients avoid damaging their fragile teeth and gums.

#### 7. Collaboration with Medical Providers:


A multidisciplinary approach is often necessary to manage the dental and systemic care of a patient with OI. Coordination with the patient's physician, physical therapist, and other specialists ensures that dental treatments are aligned with their overall health plan, particularly if there are bone deformities or other complications.

## **8. Genetic Counselling:**

Because OI is inherited, genetic counselling may be helpful for individuals with OI who are considering having children. It can help them understand the risks and implications of passing the condition on to offspring.

## **Conclusion:**

Dentists caring for patients with osteogenesis imperfecta need to be aware of the unique challenges associated with the condition, particularly in terms of tooth fragility, dental development, and overall bone health. A comprehensive, preventive approach to care, along with the use of protective measures, is key to improving the oral health and quality of life of individuals with OI. The role of both the dentist and the medical team is essential in managing the dental implications of this genetic disorder.



## References

- Huber, M. A. (2007). Osteogenesis imperfecta. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontology*, 103(3), 314-320.
- Oelerich, O., Kleinheinz, J., Böhner, L., Wiesmueller, V., & Hanisch, M. (2022). Dental implants in people with osteogenesis imperfecta: a systematic review. *International Journal of Environmental Research and Public Health*, 19(3), 1563.
- Teixeira, C. S., Felipe, M. C. S., Felipe, W. T., Silva-Sousa, Y. T. C., & Sousa-Neto, M. D. (2008). The role of dentists in diagnosing osteogenesis imperfecta in patients with dentinogenesis imperfecta. *The Journal of the American Dental Association*, 139(7), 906-914.